

Antibacterial Activity of Chlorhexidine After Final Irrigation With Ethanol: CLSM and Culture-Based Method Analysis

JULIANA MARTINS SUMAN,¹ MILTON CARLOS KUGA,² RICARDO ABREU DA ROSA,¹ MANUELA FAVARIN SANTINI,¹ RENATA GRAZZIOTIN-SOARES,¹ FRANCISCO MONTAGNER,¹ AND MARCUS VINÍCIUS REIS SÓ^{1*}

¹Department of Conservative Dentistry, Federal University of Rio Grande Do Sul, Porto Alegre, Rio Grande do Sul, Brazil

²Department of Restorative Dentistry, Araraquara Dental School, São Paulo State University, Araraquara, São Paulo, Brazil

KEY WORDS biofilms; confocal microscopy; root canal irrigants

ABSTRACT This study investigated the effect of 95% ethanol on the antibacterial properties of 2% chlorhexidine (CHX) over monospecies biofilm (*Enterococcus faecalis*) through a culture-based method, and over multispecies biofilm using confocal laser scanning microscopy (CLSM). For monospecies model, *E. faecalis* biofilm was induced in 40 root canals. The irrigation procedures were: S—saline solution; S/CHX—saline solution + CHX; E—ethanol; and E/CHX—ethanol + CHX. Microbial sampling was performed at three periods: before (S1), immediately after (S2), and 72 h after the final flush (S3). For multispecies biofilm model, 28 sterilized bovine dentin blocks were fixed on a removable orthodontic device to allow intraoral biofilm development. Seven samples were used in each group. Statistical analysis was carried out by using the Kruskal–Wallis test and Dunn’s test for multiple comparisons. There was a significant reduction in CFUs count immediately after the final flush (S2) in all experimental groups ($P < 0.05$). However, only S/CHX, E and E/CHX groups had CFU counts close to zero, without differences among them ($P > 0.05$). After 72h (S3), the S/CHX and E/CHX groups had CFU counts near zero ($P > 0.05$). The CFU count increased in S3 for S and E groups ($P < 0.05$). CLSM showed that the percentages of remaining live cells were similar in S/CHX, E, and E/CHX groups ($P > 0.05$). The S group had the highest percentage of live cells ($P < 0.05$). The 95% ethanol did not interfere in the antibacterial properties of 2% CHX over mono- and multispecies biofilms. *Microsc. Res. Tech.* 78:682–687, 2015. © 2015 Wiley Periodicals, Inc.

INTRODUCTION

In recent years, several substances have been investigated in endodontics for their use as final irrigants. Final irrigation attempts to optimize the irrigation provided by the main irrigant, and it is performed based on the associated use of two or more substances, after completing the root canal preparation (Alves et al., 2011; Baca et al., 2011; Paiva et al., 2012; Zehnder, 2006). These final irrigating substances aim, for example, to remove the smear layer (Zehnder, 2006), to dissolve remaining organic or inorganic tissue (Haapasalo et al., 2014), to reduce surface tension (Zehnder et al., 2005), and to enhance the antibacterial efficacy of the root canal treatment (Zehnder 2006). Alves et al. (2011) suggest that there may be a benefit of using passive ultrasonic irrigation (PUI) for the activation of sodium hypochlorite (NaOCl) followed by a final rinse with chlorhexidine (CHX) as supplementary steps in the treatment of infected oval-shaped root canals. Therefore, the combination of CHX, used as a final irrigant, with NaOCl, the main irrigating substance, provides benefits from both solutions.

On the other hand, authors have reported that when NaOCl and CHX come into contact, by-products might form (Prado et al., 2013; Rasimick et al., 2008). These by-products can be solid precipitates, in the form of a

brownish-reddish mass (Zehnder, 2006), that are difficult to remove, and may occlude the dentinal tubules; they can also be toxic to the periapical tissues (Akisue et al., 2010; Prado et al., 2013; Vivacqua-Gomes et al., 2002).

To cope with this problem, some substances have been suggested as intermediate irrigants to wash out the canal between the main and final irrigants (Prado et al., 2013). Although some studies have investigated the ability of intermediate irrigants to avoid precipitate formation, there is a lack of consistent evidence regarding the properties that might result from the combination of intermediate and final irrigating substances. One substance that may be used as intermediate irrigant is 95% ethanol (Krishnamurthy and Sudhakaran, 2010; Magro et al., 2015). Alcohol should be used as a flush between NaOCl and CHX, and it is indicated to remove residual NaOCl, preventing the formation of precipitates (Krishnamurthy and

*Correspondence to: Marcus Vinícius Reis SÓ, Faculty of Dentistry, Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil.
E-mail: endo-so@hotmail.com

Received 7 January 2015; accepted in revised form 1 May 2015

REVIEW EDITOR: Chuanbin Mao

DOI 10.1002/jemt.22525

Published online 3 July 2015 in Wiley Online Library (wileyonlinelibrary.com).

Sudhakaran, 2010). Nevertheless, many factors regarding the intracanal use of 95% ethanol are unknown, such as its influence on the antibacterial properties of CHX.

Therefore, the aim of this study was to investigate the effects of 95% ethanol on the antibacterial properties of CHX over monospecies biofilm (*Enterococcus faecalis*) using a culture-based method and over multispecies biofilm using confocal laser scanning microscopy (CLSM). The null hypothesis tested was that all irrigating procedures will promote similar antibacterial effect over mono- and multispecies biofilm models.

MATERIALS AND METHODS

First, a monospecies biofilm model was used and extracted teeth were inoculated with *Enterococcus faecalis* (ATCC 29212). The second method induced intraorally a multispecies biofilm. The data obtained over mono- and multispecies biofilms were assessed by using colony forming units (CFUs) count and CLSM, respectively. The sample size was based on previous report from the literature [Magro et al. (2015) for the monospecies biofilm model; Del Carpio-Perochena et al. (2011), for the multispecies biofilm method], with a 95% level of confidence and power of the study equal to 80%. The study protocol was approved by the Ethics Committees in Research of the Federal University of Rio Grande do Sul (protocol no. 317.826).

Monospecies Biofilm Model and CFU Count

Forty human mandibular premolars were selected, cleaned, sterilized, and stored in 0.9% saline solution at 4°C. All teeth were radiographed to confirm the presence of one canal, the absence of calcification, resorptions, immature foramen, and root canal filling. The crowns were removed and the working length was standardized at 12 mm, 1 mm shorter than the apical foramen as visually determined.

In order to standardize the width of the canals, they were instrumented under constant irrigation/aspiration with 2.5% NaOCl using the ProTaper Universal® system until F2 instrument. After preparation, the canals were irrigated with 17% EDTA (Biodinâmica, Iporã, PR, Brazil) and then irrigated with 5 mL of saline to wash out the EDTA. The apices were then sealed with composite resin (Magicfill Vigodent-Coltene, São Paulo, SP, Brazil) and the roots were sterilized using ethylene oxide. The efficacy of the sterilization was verified by inserting paper points into the canals, putting them into polypropylene tubes (Eppendorf do Brasil Ltda, São Paulo, SP, Brazil), and subsequently maintaining the tubes in an incubator at 37°C for 7 days. The solution turbidity was evaluated every 24 h.

The microbiological procedures were performed under aseptic conditions, in a laminar flow chamber. *E. faecalis* strain was used to infect the canals. A suspension was prepared by adding 1 mL of a pure culture of *E. faecalis* grown in brain–heart infusion broth (BHI; Difco, Detroit, MI) for 24 h at 37°C in a bacteriological incubator. The concentration of the obtained microbial suspension was adjusted to 0.5 on the McFarland scale. Fifty microliters of the microbial broth was inoculated inside the canal and the roots were maintained for 7 days at 37°C in the incubator.

Every two days, 25 µL of the microbial broth was removed from the root canals and replaced for fresh BHI broth q.s.p. to ensure cell nutrition and viability.

Microbial collection and counting were performed at three time points: before (S1), immediately after (S2), and 72 h after the final flush (S3). The initial sample (S1) was performed using three size 25 sterile paper points (Tanariman, Manacapuru, AM, Brazil) placed inside the root canal for 1 min and then transferred to a polypropylene tube (Eppendorf) containing 1 mL of sterile saline solution. The tubes were vortexed for 1 min (Vortex Ap 56; Phoenix, Araraquara, SP, Brazil). After decimal serial dilutions, aliquots were seeded into Petri dishes containing BHI agar (Difco) and incubated at 37°C for 24 h to allow the counting of CFU. The culture purity was confirmed by the colony morphology, Gram staining, and the catalase test. No sample was discarded due to absence of culture purity.

After that, the irrigation procedures were performed as follows: each irrigant (5 mL) was inserted into the root canal using a plastic syringe equipped with a NaviTip® needle (Ultradent Products, Inc., South Jordan, UT). Once the irrigation was complete, the substance was then aspirated with an endodontic cannula. The four treatments used were: S—saline; S/CHX—saline (5 min) + CHX (5 min); E—ethanol (5 min); and E/CHX—ethanol (5 min) + CHX (5 min). After the irrigation procedures, S2 and decimal serial dilutions were performed identically to S1.

All canals were filled with 50 µL of BHI broth after S2 and the roots were stored at 37°C. After 72 h, S3 was performed as described before. One evaluator blinded to the experimental groups performed the microbial counting always 24 h after the sample collections.

The intergroup analysis was assessed using Kruskal–Wallis and Dunn's tests to compare the CFUs' count after each microbial collection. Intragroup assessment was performed using Friedman's test to compare each protocol at different time points. The significance level was set at 5% for all statistical tests.

Multispecies Biofilm Method and CLSM Analysis

Twenty eight sterile bovine dentin blocks (3 mm × 3 mm × 2 mm) were used. The samples were treated with 17% EDTA for 3 min to eliminate the smear layer produced during the sectioning process.

All dentin blocks were fixed on a removable orthodontic device to allow intraoral biofilm development (Del Carpio-Perochena et al., 2011). After signing an informed consent form, two healthy volunteers (aged 26 and 29) who had similar eating habits used the intraoral device continuously for 72 h. Both volunteers only removed the intraoral device when consuming food and drink and when practicing oral hygiene. At the end of the infection period, the dentin blocks were removed from the orthodontic device and inserted into a polypropylene tube with 1 mL of BHI (Difco). The tubes were then stored in an incubator at 37°C (Ordinola-Zapata et al., 2013).

After 24 h, the dentin blocks were removed from the tubes and then rinsed with 1 mL of distilled water to remove the BHI medium and any non-adherent cells. Next, they were randomly divided in four groups ($n = 7$): S—saline (5 min); S/CHX—saline (5

min) + CHX (5 min); E—ethanol (5 min); and E/CHX—ethanol (5 min) + CHX (5 minutes). Each specimen was immersed in 5 mL of the irrigant.

The analysis of biofilm viability was performed by using the SYTO 9/propidium iodide technique (Live/Dead, BacLight, Invitrogen, Eugene, OR). SYTO 9 is a green fluorescent stain that labels both live and dead microorganisms. Propidium iodide is a red fluorescent nucleic acid stain that only penetrates cells with damaged membranes (dead cells). First, the samples were cleaned with 2 mL of saline solution and then 0.25 μ L of dye was placed over the biofilm. A CLSM (Olympus Fluoview 1000, Olympus Corporation, Tokyo, Japan) was used to visualize the samples (Ordinola-Zapata et al., 2012). The respective absorption and emission wavelengths were 494/518 nm for SYTO 9 and 536/617 nm for propidium iodide. The biofilm was randomly assessed at $\times 60$ magnification. Next, three confocal stacks from different random areas were obtained from each sample using a $\times 40$ oil lens, a 1- μ m step size, and a format of 386×386 pixels. A total number of 21 stacks (3 operative fields \times 7 specimens per group) were obtained for each group. The evaluator was blinded to the experimental groups. All images

were analyzed using BioImage_L software (<http://bioimage.com>) for the total biovolume (μm^3), the total number of live cells (green), and the percentage of live cells (Chavez de Paz, 2009). The percentage of live cells (green) was assessed by using the Kruskal–Wallis test and Dunn's test for multiple comparisons ($\alpha = 5\%$).

RESULTS

In the monospecies biofilm model, there was a significant reduction in CFUs count immediately after the final flush (S2) for all experimental groups ($P < 0.05$) (Table 1). However, only the S/CHX, E and E/CHX groups presented counts near zero with no differences between them ($P > 0.05$). After 72 h (S3), the S/CHX and E/CHX groups maintained CFUs' count near zero ($P > 0.05$) (Table 2). The CFUs' count increased significantly in S3 for S and E groups ($P < 0.05$).

CLSM showed that the percentages of remaining live cells were similar in the S/CHX, E, and E/CHX groups ($P > 0.05$). The S group presented the highest percentage of live cells ($P < 0.05$) (Table 3). Table 3 also presents the total biovolume ($\mu\text{m}^3 \times 10^6$), the biovolume of live cells, and the percentage of live cells (multispecies biofilm). Figure 1 presents CLSM representative images of multispecies biofilms after the four irrigation procedures.

DISCUSSION

Alcohol has been indicated for use as an intermediate flush between NaOCl (main irrigant) and CHX (final irrigant) because it can prevent the formation of precipitates that occur as a consequence of mixing the two substances (Krishnamurthy and Sudhakaran, 2010). As reported previously, isopropyl alcohol prevented the formation of the insoluble neutral salt that results from the interaction of NaOCl and CHX (Krishnamurthy and Sudhakaran, 2010). Short et al. (2003) showed that 96% ethyl alcohol and 70% isopropyl alcohol were efficient in removing NaOCl crystals on the gutta-percha cones after immersion in 5.25% and 2.5% NaOCl for 1 min (Short et al., 2003). Thus, despite a recent study showed opposite results (Magro et al., 2015), 95% alcohol still seems adequate to remove residual NaOCl to prevent the formation of the precipitate (Krishnamurthy and Sudhakaran, 2010; Short et al., 2003). It is probable that the dilution of absolute alcohol with distilled water may compromise their ability to remove the precipitates formed from the interaction between the NaOCl and CHX, despite the known antimicrobial activity of 70% alcohol (Magro et al., 2015).

Therefore, this study evaluated the influence of 95% ethanol on the antibacterial properties of CHX. The present results showed that the use of 95% ethanol as

TABLE 1. Total percentage of remaining live cells after the microbial collection from the root canals in each group according to the experimental period: before final flush (S1), immediately after final flush (S2), and after 72 h (S3)

Irrigants	Experimental periods		
	S1	S2	S3
Saline	100% ^b	9.5% ^{Ba}	33.9% ^{Bb}
Saline + CHX	100% ^b	0% ^{Aa}	0% ^{Aa}
Ethanol	100% ^b	0% ^{Aa}	34.3% ^{Bb}
Ethanol + CHX	100% ^b	0.1% ^{Aa}	0% ^{Aa}

CHX, chlorhexidine.

Different superscript capital letters in each column represent statistical significance ($P < 0.05$), Kruskal–Wallis test and Dunn test ($\alpha = 5\%$).

Different superscript lower case letters in each row represent statistical significance ($P < 0.05$), Friedman's test ($\alpha = 5\%$).

TABLE 2. Comparison of CFUs between groups in each experimental period: before final flush (S1), immediately after final flush (S2), and after 72 h (S3)

Irrigants	S1	S2	S3
Saline	160 ^B	16.4 ^{Ab}	21.2 ^{ABb}
Saline + CHX	306 ^B	0.0125 ^{Aa}	0.0104 ^{Aa}
Ethanol	40 ^B	0.0009 ^{Aa}	22 ^{Bb}
Ethanol + CHX	128 ^B	0.8 ^{Aa}	0.01 ^{Aa}

CHX, chlorhexidine.

Different superscript capital letters in each row represent statistical significance ($P < 0.05$), Friedman's test ($\alpha = 5\%$).

Different superscript lowercase letters in each column represent differences ($P < 0.05$) after Kruskal–Wallis and Dunn post hoc tests ($\alpha = 5\%$).

TABLE 3. Median and percentiles P(25) and P(75) of cell biovolume and percentage of live cells on bovine dentine samples in each group

		Saline (n = 7)	Saline + CHX (n = 7)	Ethanol (n = 7)	Ethanol + CHX (n = 7)
Median, P(25), P(75)	Total biovolume ($\mu\text{m}^3 \times 10^6$)	16 (8.4/17.2)	21.2 (16.9/28.2)	21.4 (15.4/29.8)	15.9 (15.1/21.6)
	Biovolume of live cells ($\mu\text{m}^3 \times 10^6$)	15 (7.3/16.7)	13.7 (12.5/15.8)	13.8 (8.6/20.3)	13.8 (12.7/15.7)
	Live cells ^a (%)	93.4 (92.7/96.5)	79.6 (65.9/81.0) ^b	64.6 (53.3/75.7) ^b	76.3 (68.9/83.5) ^b

CHX, chlorhexidine.

^a(Biovolume of live cells/Total biovolume) \times 100.

^bStatistical significant difference in relation to saline group, Kruskal–Wallis test and Dunn test ($\alpha = 5\%$).




Fig. 1. CLSM representative images of multispecies biofilm model after the four irrigation procedures: (A) saline; (B) saline/chlorhexidine; (C) ethanol; and (D) ethanol/chlorhexidine. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

an intermediate irrigating substance did not affect the antibacterial properties of CHX over mono- and multispecies biofilms (Tables (1–3)). The groups S and E received half the total volume of irrigation when compared to the groups S/CHX and E/CHX. There was a significant decrease in the microbial load for all the groups from S1 to S2, which can be associated with the mechanical effect of the irrigation. The increase of the CFU values at S3 was observed for groups S and E. The presence of residual microorganisms inside the root canal system and the limited antimicrobial actions of S and E may explain this behavior.

In the current study, results of viability for multispecies biofilm can be compared with those obtained at S2 in monospecies biofilm model because both assessed the antibacterial effect of the experimental protocols immediately after the contact of the microorganisms with the irrigants. In this scenario, both methods showed similar results. UFCs' count and CLSM assessments indicated that E, CHX, and E/CHX groups presented similar antibacterial properties ($P > .05$). The antibacterial effect of CHX has been well established in the literature (Du et al., 2014; Stojicic et al., 2013). Figure 1 shows a number of viable cells incompatible with Table 2, especially in the Saline + CHX and E + CHX groups. It is important to understand that when working with CSLM some images are captured in a deeper region of the biofilm where the chemical was not effective (Fig. 2). It should be recognized that no in vitro method can accurately reflect the conditions under which the microorganisms grow and develop within the root canal system. The multispecies biofilm model take into account that the endodontic pathogens are less susceptible to the action of antimicrobial agents when organized in communities adhered to the




Fig. 2. Images from CLSM indicating predominance of live cells in deeper layers of biofilms (A, C, and E). In contrast, in biofilm superficial layers the irrigants were more effective in killing microorganisms (B, D, and F). A, B—Saline/CHX; C, D—ethanol; and E/F—ethanol/CHX. [Color figure can be viewed in the online issue, which is available at wileyonlinelibrary.com.]

walls of the root canal. It can be one of the main reasons for the persistence of the microorganisms after disinfection protocols.

Another relevant finding from the present study was that 95% ethanol, when used alone, had antibacterial effects against mono- and multispecies biofilms. It is well-known that alcohol promotes disinfection of gutta-percha cones (Short et al., 2003). One explanation for this finding is that alcohol (both ethyl alcohol and isopropyl alcohol) is a tensioactive agent; it is highly electronegative (Krishnamurthy and Sudhakaran, 2010). Due to the low surface tension and the physical action of the irrigation procedure (irrigation/aspiration), 95% ethanol significantly reduced the

microbial count in the root canal at S2 (monospecies biofilm) and reduced the percentage of live cells (multispecies biofilm). On the other hand, Prado et al. (2013) observed that the association E/CHX produced a salt precipitation after 5 s. The authors attributed the precipitate formation to the reduced solubility of the CHX salt in ethanol.

Microorganisms organized in biofilms may become established on any organic or inorganic surface substrate, and planktonic microorganisms prevail in a water-based solution (Mohammadi and Abbott, 2009). The literature has proven that when microorganisms are organized in biofilms, they are more resistant than the corresponding planktonic form of the same microorganism (Mohammadi and Abbott, 2009; Svensater and Bergenholtz, 2004). In this regard, the present study used both forms of microorganism organization. Therefore, mono- and multispecies biofilms were induced and assessed using two different methodologies. Culture-based methods are useful due to their relative practicality and reduced cost. In addition to these advantages, they permit the ability to visually count the microorganisms; i.e., under ideal conditions of plating and incubation, the cells (or small cell clusters) grow singly, resulting in colonies known as CFUs (Fonseca et al., 2008). The use of CLSM is valuable for studying the structure of biofilms because it allows investigation of the intact biofilm without the interference of sample processing (Hohscheidt et al., 2013; Shen et al., 2009; Zapata et al., 2008). The immersion protocol employed to remove the multispecies biofilm formed over the dentin blocks did not allow to reproduce the dynamics of the irrigants that occur inside the root canal system. However, the aim of the study was to evaluate the antimicrobial properties of the irrigants alone or in association, and its potential residual effect.

The multispecies biofilm formation was induced in a manner that was very similar to real clinical conditions. On the other hand, although the use of a monospecies biofilm (*E. faecalis* inoculated in the root canals) was useful for investigating the current hypothesis, the use of a planktonic polymicrobial infection is suggested for further studies. Regarding the microbial counting performed at S3, this study showed that there was inhibition of microbial growth in the two groups with CHX (S/CHX and E/CHX). This inhibition probably occurred due to the residual effectiveness of the CHX, which is in line with the work by Baca et al. (2011). These authors incubated *E. faecalis* in extracted teeth to evaluate the antimicrobial effects of different final irrigation procedures. They concluded that an effective final irrigation regimen would be a combination of 2% CHX + 0.2% Cetrimide (a cationic surfactant agent), which exerts antimicrobial action over time.

Despite this advantage, the biocompatibility of ethanol when used as an endodontic irrigant is not yet well established (Krishnamurthy and Sudhakaran, 2010) and its effects on the periapical tissues remain a concern. Therefore, the results of the present study may not be directly applied as a clinical protocol. According to the results of the present study, 95% ethanol did not interfere in the antibacterial properties of 2% CHX solution over mono- and multispecies biofilms.

ACKNOWLEDGMENT

The authors deny any conflicts of interest related to this study.

REFERENCES

- Akisue E, Tomita VS, Gavini G, Poli de Figueiredo JA. 2010. Effect of the combination of sodium hypochlorite and chlorhexidine on dentinal permeability and scanning electron microscopy precipitate observation. *J Endod* 36:847–850.
- Alves FR, Almeida BM, Neves MA, Moreno JO, Rôças IN, Siqueira JF, Jr. 2011. Disinfecting oval-shaped root canals: Effectiveness of different supplementary approaches. *J Endod* 37:496–501.
- Baca P, Mendoza-Llamas ML, Arias-Moliz MT, González-Rodríguez MP, Ferrer-Luque CM. 2011. Residual effectiveness of final irrigation regimens on enterococcus faecalis-infected root canals. *J Endod* 37:1121–1123.
- Chavez de Paz LE. 2009. Image analysis software based on color segmentation for characterization of viability and physiological activity of biofilms. *Appl Environ Microbiol* 75:1734–1739.
- Del Carpio-Perochena AE, Bramante CM, Duarte MA, Cavenago BC, Villas-Boas MH, Graeff MS, Bernardineli N, de Andrade FB, Ordinola-Zapata R. 2011. Biofilm dissolution and cleaning ability of different irrigant solutions on intraorally infected dentin. *J Endod* 37:1134–1138.
- Du T, Wang Z, Shen Y, Ma J, Cao Y, Haapasalo M. 2014. Effect of long-term exposure to endodontic disinfecting solutions on young and old enterococcus faecalis biofilms in dentin canals. *J Endod* 40:509–514.
- Fonseca MB, Júnior PO, Pallota RC, Filho HF, Denardin OV, Rapoport A, Dedivitis RA, Veronezi JF, Genovese WJ, Ricardo AL. 2008. Photodynamic therapy for root canals infected with enterococcus faecalis. *Photomed Laser Surg* 26:209–213.
- Haapasalo M, Shen Y, Wang Z, Gao Y. 2014. Irrigation in endodontics. *Br Dent J* 216:299–303.
- Hohscheidt GL, Böttcher DE, Fatturi Parolo CC, Montagner F, Grecca FS. 2013. Response of *E. faecalis* biofilms to different associations of auxiliary substances during root canal preparation: A confocal laser microscopy analysis. *Microsc Res Tech* 76:658–662.
- Krishnamurthy S, Sudhakaran S. 2010. Evaluation and prevention of the precipitate formed on interaction between sodium hypochlorite and chlorhexidine. *J Endod* 36:1154–1157.
- Magro MG, Kuga MC, Aranda-Garcia AJ, Victorino KR, Chávez-Andrade GM, Faria G, Keine KC, Só MV. 2015. Effectiveness of several solutions to prevent the formation of precipitate due to the interaction between sodium hypochlorite and chlorhexidine and its effect on bond strength of an epoxy-based sealer. *Int Endod J* 48:478–483.
- Mohammadi Z, Abbott PV. 2009. The properties and applications of chlorhexidine in endodontics. *Int Endod J* 42:288–302.
- Ordinola-Zapata R, Bramante CM, Cavenago B, Graeff MS, Gomes de Moraes I, Marciano M, Duarte MA. 2012. Antimicrobial effect of endodontic solutions used as final irrigants on a dentine biofilm model. *Int Endod J* 45:162–168.
- Ordinola-Zapata R, Bramante CM, Minotti PG, Cavenago BC, Garcia RB, Bernardineli N, Jaramillo DE, Hungaro Duarte MA. 2013. Antimicrobial activity of triantibiotic paste, 2% chlorhexidine gel, and calcium hydroxide on an intraoral-infected dentin biofilm model. *J Endod* 39:115–118.
- Paiva SS, Siqueira JF, Jr, Rôças IN, Carmo FL, Ferreira DC, Curvelo JA, Curvelo JA, Soares RM, Rosado AS. 2012. Supplementing the antimicrobial effects of chemomechanical debridement with either passive ultrasonic irrigation or a final rinse with chlorhexidine: A clinical study. *J Endod* 38:1202–1206.
- Prado M, Santos Júnior HM, Rezende CM, Pinto AC, Faria RB, Simão RA, Gomes BP. 2013. Interactions between irrigants commonly used in endodontic practice: A chemical analysis. *J Endod* 39:505–510.
- Rasimick BJ, Nekich M, Hladek MM. 2008. Interaction between chlorhexidine gluconate and EDTA. *J Endod* 34:1521–1523.
- Shen Y, Qian W, Chung C, Olsen I, Haapasalo M. 2009. Evaluation of the effect of two chlorhexidine preparations on biofilm bacteria in vitro: A three-dimensional quantitative analysis. *J Endod* 35:981–985.
- Short RD, Dorn SO, Kuttler S. 2003. The crystallization of sodium hypochlorite on gutta-percha cones after the rapid-sterilization technique: An SEM study. *J Endod* 29:670–673.
- Stojicic S, Shen Y, Haapasalo M. 2013. Effect of the source of biofilm bacteria, level of biofilm maturation, and type of disinfecting agent

- on the susceptibility of biofilm bacteria to antibacterial agents. *J Endod* 39:473–477.
- Svensater G, Bergenholtz G. 2004. Biofilms in endodontic infections. *Endod Topics* 9:27–36.
- Vivacqua-Gomes N, Ferraz CC, Gomes BP, Zaia AA, Teixeira FB, Souza-Filho FJ. 2002. Influence of irrigants on the coronal microleakage of laterally condensed gutta-percha root fillings. *Int Endod J* 35:791–795.
- Zapata RO, Bramante CM, de Moraes IG, Bernardineli N, Gasparoto TH, Graeff MS, Campanelli AP, Garcia RB. 2008. Confocal laser scanning microscopy is appropriate to detect viability of enterococcus faecalis in infected dentin. *J Endod* 34:1198–1201.
- Zehnder M. 2006. Root canal irrigants. *J Endod* 32:389–398.
- Zehnder M, Schicht O, Sener B, Schmidlin P. 2005. Reducing surface tension in endodontic chelator solutions has no effect on their ability to remove calcium from instrumented root canals. *J Endod* 31:590–592.